



TITLE:

On the Cohomology of the Moduli space of Parabolic Connections

AUTHOR(S):

Matsubara, Yuki

CITATION:

Matsubara, Yuki. On the Cohomology of the Moduli space of Parabolic Connections. 代数幾何学シンポジウム記録 2018, 2018: 147-147

ISSUE DATE:

2018

URL:

<http://hdl.handle.net/2433/236415>

RIGHT:

ON THE COHOMOLOGY OF THE MODULI SPACE OF PARABOLIC CONNECTIONS

Yuki Matsubara (D2)

Department of Mathematics, Graduate School of Science, Kobe University

Introduction

In this poster, we study the moduli space of logarithmic connections of rank 2 on $\mathbb{P}^1 \setminus \{t_1, \dots, t_5\}$ with fixed spectral data. We compute the cohomology of such moduli space, and this computation will be used to extend the results of geometric Langlands correspondence of [1] to the case where the parabolic connections have five simple poles on \mathbb{P}^1 .

Preliminaries

We introduce \mathfrak{sl}_2 -connections.

Fix complex numbers $\nu_1, \dots, \nu_n \in \mathbb{C}$. Suppose that $\nu_1 \cdots \nu_n \neq 0$ and

$$\sum_{i=1}^n \epsilon_i \nu_i \notin \mathbb{Z} \quad (1)$$

for any $(\epsilon_i), \epsilon_i \in \{1, -1\}$.

Definition 2.1 A ν - \mathfrak{sl}_2 -parabolic connection on \mathbb{P}^1 is a triplet (E, ∇, φ) such that

1. E is a rank 2 vector bundle on \mathbb{P}^1 ,
2. $\nabla : E \rightarrow E \otimes \Omega_{\mathbb{P}^1}^1(D)$ is a connection, where $D := t_1 + \cdots + t_n$,
3. $\varphi : \wedge^2 E \simeq \mathcal{O}_{\mathbb{P}^1}(-1)$ is a horizontal isomorphism,
4. the residue $\text{res}_i(\nabla)$ of the connection ∇ at t_i has eigenvalues ν_i^{\pm} , $1 \leq i \leq n$.

Here, we put

$$\nu_i^{\pm} := \pm \nu_i \quad (i = 1, \dots, n-1), \quad \nu_n^+ := \nu_n, \quad \nu_n^- := 1 - \nu_n.$$

Denote by \mathcal{M} the moduli stack of ν - \mathfrak{sl}_2 -parabolic connections on \mathbb{P}^1 , and by M its coarse moduli space.

For such ν , the parabolic direction $l_i := \ker(\text{res}_i(\nabla) - \nu_i) \subset E|_{t_i}$ is uniquely determined. So, we can get the forgetful map

$$\text{Bun} : M \rightarrow P := (E, \nabla, \varphi) \mapsto (E, l_i)$$

where P is the coarse moduli space of undecomposable quasi-parabolic bundles (E, l_i) on \mathbb{P}^1 .

Now, we consider the following stratification of M . By the irreducibility of $(E, \nabla, \varphi) \in M$, we have the following proposition.

Proposition 2.2 For $(E, \nabla, \varphi) \in M$, we have

$$E \simeq \mathcal{O}(k) \oplus \mathcal{O}(-k-1) \quad \text{where } 0 \leq k \leq \left\lfloor \frac{n-3}{2} \right\rfloor.$$

Denote by M^k the subvariety of M where $E \simeq \mathcal{O}(k) \oplus \mathcal{O}(-k-1)$. Then

$$M = M^0 \cup \cdots \cup M^{\lfloor (n-3)/2 \rfloor}.$$

Note that the stratum M^0 is a Zariski open dense of M .

Geometric description of M^0

Suppose $n = 5$. For computation of the cohomology of M , we introduce some blowing-up of the Hirzebruch surface \mathbb{F}_3 . Put $L := \mathcal{O}_{\mathbb{P}^1}^*(D)$. Let \tilde{L} be the total space of the line bundle L . Note that $\tilde{L} = \mathbb{F}_3 \setminus s_{\infty}$ where s_{∞} is the infinity section ($s_{\infty}^2 = -3$). Let $\pi : \tilde{L} \rightarrow \mathbb{P}^1$ be the projection and let $\tau_i : \tilde{F}_i \simeq \pi^{-1}(t_i) \xrightarrow{\sim} \mathbb{C}$ be the residue map. Put $\nu_i^+ := \nu_i$, $\nu_i^- := -\nu_i$ for $i = 1, \dots, 4$, $\nu_5^+ := \nu_5$, $\nu_5^- := 1 - \nu_5$ and $\hat{\nu}_i^{\pm} := \tau_i^{-1}(\nu_i^{\pm})$. Set

$$K_{5,q}^{\pm} := (\text{Bl}_{\nu^{\pm}} \tilde{L}) \setminus (\tilde{F}_1 \cup \cdots \cup \tilde{F}_5)$$

where $\text{Bl}_{\nu^{\pm}} \tilde{L}$ is the blowing-up of \tilde{L} at $\hat{\nu}_i^{\pm}$ for $i = 1, \dots, n$, and \tilde{F}_i are the proper pre-images of the fiber F_i $i = 1, \dots, n$.

Lemma 3.1

$$H^i(K_{5,q}^{\pm}, \mathcal{O}_{K_{5,q}^{\pm}}) = \begin{cases} \mathbb{C}, & i = 0, \\ H_n^{\pm}(A) \neq 0, & i = 1, \\ 0, & i \geq 2, \end{cases}$$

where (A, \mathfrak{m}) is a local ring such that $\dim(A_{\mathfrak{m}}) = 2$.

We denote by K_5 the image of $K_{5,q}^{\pm}$ under the projection $K_{5,q}^{\pm} \rightarrow \mathbb{F}_3 \setminus s_{\infty}$. We can define the following map

$$M^0 \rightarrow \text{Sym}^2(K_5) \\ (E, \nabla, \varphi) \mapsto \{(q_1, p_1), (q_2, p_2)\}, \quad (2)$$

We consider the composite of the Hilbert-Chow morphism and the blowing-up

$$\text{Hilb}^2(K_5) \rightarrow \text{Sym}^2(K_5) \rightarrow \text{Sym}^2(K_5).$$

We have the next proposition.

Proposition 3.2 ([3] Theorem 5.2) We can extend the map (2) to

$$M^0 \longrightarrow \text{Hilb}^2(K_5)$$

and this map is injective.

Cohomology of $K_{5,q}^{\pm}$

We denote by $Z \subset \text{Sym}^2(K_5)$ the proper pre-image of $\{(q_1, p_1), (q_1, -p_1)\} \subset \text{Sym}^2(K_5)$ under the blowing-up $\text{Sym}^2(K_5) \rightarrow \text{Sym}^2(K_5)$, and by $\tilde{Z} \subset \text{Hilb}^2(K_5)$ the proper pre-image of Z under the Hilbert-Chow morphism $\text{Hilb}^2(K_5) \rightarrow \text{Sym}^2(K_5)$. Denote by

$$\widetilde{\text{Hilb}}^2(K_5) \rightarrow \text{Hilb}^2(K_5) \quad (3)$$

the blowing-up along \tilde{Z} and by \tilde{Z} the strict transform of \tilde{Z} . We also denote by $(K_5^{\pm} \times K_5^{\pm})^{\sim}$ the blowing-up of $K_5^{\pm} \times K_5^{\pm}$ along the ideal $(q_1 - q_2, p_1 - p_2)$, and by $(K_5^{\pm} \times K_5^{\pm})^{\sim}$ the blowing-up of $(K_5^{\pm} \times K_5^{\pm})^{\sim}$ along the ideal $(q_1 - q_2, p_1 + p_2)$. Then $\text{Hilb}^2(K_5^{\pm}) = (K_5^{\pm} \times K_5^{\pm})^{\sim} / \mathfrak{S}_2$ and $\widetilde{\text{Hilb}}^2(K_5^{\pm}) = (K_5^{\pm} \times K_5^{\pm})^{\sim} / \mathfrak{S}_2$. Now, using above description, we define another important blowing-up of Hirzebruch surface \mathbb{F}_3 . Fix $q_i \in \mathbb{P}^1 \setminus \{t_1, \dots, t_5\}$ and define the fiber \tilde{F}_i over q_i . We denote by $(\mathbb{F}_3)^{\sim}$ the blowing-up of \mathbb{F}_3 at two points $\{(q_1, p_1), (q_1, -p_1)\}$ (when $p_1 = p_2 = 0$, blow up twice at $(q_1, 0)$). Set

$$K_{5,q_1}^{\pm} := (\mathbb{F}_3)^{\sim} \setminus (s_{\infty} \cup \tilde{F}_1 \cup \cdots \cup \tilde{F}_5)$$

where \tilde{F}_i is the strict transform of F_i . We denote by K_{5,q_1} the image of K_{5,q_1}^{\pm} under the projection $K_{5,q_1}^{\pm} \rightarrow \mathbb{F}_3 \setminus s_{\infty}$. Define

$$D_{q_1} := 2s_{\infty} + \tilde{F}_1 + \cdots + \tilde{F}_5 + \tilde{F}_5$$


Then, we have

$$(s_{\infty}, \tilde{F}_i) = 1, (\tilde{F}_i, \tilde{F}_i) = 0, (s_{\infty}, s_{\infty}) = -3, (\tilde{F}_i, \tilde{F}_i) = -2$$

This implies

$$(D_{q_1}, D_{q_1}) = (D_{q_1}, s_{\infty}) = (D_{q_1}, \tilde{F}_i) = 0.$$

This is the same condition of Arinkin Lysenko [2], section 5.2.



Proposition 4.1

$$H^i(K_{5,q}^{\pm}, \mathcal{O}_{K_{5,q}^{\pm}}) = \begin{cases} \mathbb{C}, & i = 0, \\ 0, & i > 0. \end{cases}$$

Cohomology of M

Set $\widetilde{M}_Z := M^0 \cup \tilde{Z}$. By Proposition 3.2, we have injective maps $\iota : M^0 \hookrightarrow \text{Hilb}^2(K_5)$ and $\iota : \widetilde{M}_Z \hookrightarrow \widetilde{\text{Hilb}}^2(K_5)$. We define the blowing-up parameter $\lambda := p_1 + p_2 = \lambda - (q_1 - q_2)$. Set $T := \widetilde{\text{Hilb}}^2(K_5) \setminus \widetilde{M}_Z$. For a vector bundle \mathcal{F} on $\widetilde{\text{Hilb}}^2(K_5)$,

$$H^i(\widetilde{M}_Z, \mathcal{F}|_{\widetilde{M}_Z}) = H^i(\widetilde{\text{Hilb}}^2(K_5), \iota_* \iota^* \mathcal{F}) \\ = \varinjlim H^i(\widetilde{\text{Hilb}}^2(K_5), \mathcal{F}(kT)).$$

To compute $H^i(\widetilde{\text{Hilb}}^2(K_5), \mathcal{F}(kT))$, consider $H^i((K_5^{\pm} \times K_5^{\pm})^{\sim}, \mathcal{F}(kT^{\vee}))$, where T^{\vee} is defined by $(\lambda_{\infty} = \infty)$. We can define a map

$$f : (K_5^{\pm} \times K_5^{\pm})^{\sim} \setminus T^{\vee} \rightarrow K_5^{\pm} \\ (q_1, p_1, q_2, p_2) \mapsto (q_1, p_1),$$

and the fiber $f^{-1}(\{(q_1, p_1)\}) \simeq K_{5,q_1}^{\pm}$. By Leray's spectral sequence, we have

$$H^i((K_5^{\pm} \times K_5^{\pm})^{\sim} \setminus T^{\vee}, \mathcal{F}) \simeq \bigoplus_{p+q=i} H^p(K_{5,q_1}^{\pm}, R^q f_* \mathcal{F}).$$

Using the Base change theorem, we have $(R^q f_* \mathcal{F})_{(q_1, p_1)} \simeq H^q(K_{5,q_1}^{\pm}, \mathcal{F}|_{(q_1, p_1)})$. By using Lemma 3.1 and Proposition 4.1, we have

$$H^i((K_5^{\pm} \times K_5^{\pm})^{\sim} \setminus T^{\vee}, \mathcal{O}) = \begin{cases} \mathbb{C}, & i = 0, \\ H_n^{\pm}(A) \neq 0, & i = 1, \\ 0, & i > 1. \end{cases}$$

Moreover, the action of \mathfrak{S}_2 on $H^1((K_5^{\pm} \times K_5^{\pm})^{\sim} \setminus T^{\vee}, \mathcal{O})$ is nontrivial. Therefore,

$$H^i(\widetilde{M}_Z, \mathcal{O}_{\widetilde{M}_Z}) = \begin{cases} \mathbb{C}, & i = 0, \\ 0, & i > 0. \end{cases}$$

Since $\text{codim}_{\widetilde{\text{Hilb}}^2(K_5)}(\tilde{Z}) = 2$, and $M^1 = M \setminus M^0 \simeq A^2$, we have

Theorem 5.1

$$H^i(M, \mathcal{O}_M) = \begin{cases} \mathbb{C}, & i = 0, \\ 0, & i > 0. \end{cases}$$

References

- [1] D. Arinkin, *Orthogonality of natural sheaves on moduli stacks of $SL(2)$ -bundles with connections on P^1 minus 4 points*, Selecta Math., New Series 7 (2001), 213-239.
- [2] D. Arinkin, S. Lysenko, *On the moduli of $SL(2)$ -bundles with connections on $\mathbb{P}^1 \setminus \{x_1, \dots, x_4\}$* , Internat. Math. Res. Notices (1997), no. 19, 983-999.
- [3] A. Kuroyo, M.-H. Saito, *Explicit description of jumping phenomena on moduli spaces of parabolic connections and Hilbert schemes of points on surfaces*, accepted in Kyoto Journal of Mathematics, (arXiv:math/1611.00971)